

# Highly accurate surface and volume integration for level set methods using a moment-fitting approach

Björn Müller<sup>\*,1,2</sup>, Florian Kummer<sup>2</sup>, Martin Oberlack<sup>1,2,3</sup>

<sup>1</sup> Graduate School CE, Dolivostraße 15, 64293 Darmstadt, Germany

<sup>2</sup> Chair of Fluid Dynamics, Petersenstraße 30, 64287 Darmstadt, Germany

<sup>3</sup> Center of Smart Interfaces, Petersenstraße 17, 64287 Darmstadt, Germany

## Abstract

We present methods for the numerical integration of functions over domains that are defined by the zero iso-contour of a level set function. Such integrals appear in many contexts, but have become a major area of interest due to the recent rise of sharp-interface methods that are trying to resolve local effects with sub-cell accuracy. Examples include the eXtended Finite Element Method (XFEM), the Finite Cell Method and the Discontinuous Galerkin Method. In a nutshell, such methods share the property that part of the burden of discretization is shifted to the numerical integration of generic functions over complicated sub-domains for which conventional quadrature are not available. The performance of these methods is thus directly linked to the affordable integration accuracy.

A number of measures to cope with this issue have been proposed in literature. Typically, they are either based on a regularization of discontinuous/singular integrands (e.g., [2]) or an adaptive subdivision of cells intersected by the interface (e.g., [3, 5]). Both approaches reach second-order convergence rates at best, which, especially in case of higher order methods, is often insufficient.

Recently, a method based on the optimization of the non-linear *moment-fitting* equations

$$\begin{pmatrix} \int f_1(\mathbf{x}) d\mathbf{x} \\ \vdots \\ \int f_M(\mathbf{x}) d\mathbf{x} \end{pmatrix} = \begin{pmatrix} f_1(\mathbf{x}_1) & \dots & f_1(\mathbf{x}_N) \\ \vdots & \ddots & \vdots \\ f_M(\mathbf{x}_1) & \dots & f_M(\mathbf{x}_N) \end{pmatrix} \begin{pmatrix} w_1 \\ \vdots \\ w_N \end{pmatrix}$$

with given functions  $f_1, \dots, f_M$ , quadrature nodes  $\mathbf{x}_1, \dots, \mathbf{x}_N$  and quadrature weights  $w_1, \dots, w_N$  has been developed [1]. If the left-hand side can be obtained, it can be used for the pre-calculation of extremely efficient quadrature rules on very general domains. In [6], the authors restrict themselves to piecewise linear interfaces, simplify the left-hand side and show how the non-linear optimization process can be reduced to a linear one in this context. The proposed modifications render the approach practicable in a broader range of applications comprising moving level sets. Still, the restriction to piecewise linear interface approximations makes a refinement near curved boundaries inevitable and, if high accuracy is required, the process becomes inefficient.

The present work aims at removing the mentioned drawbacks without necessitating a complex, non-general higher order interface reconstruction. On the contrary, the presented methods allow for a realization of the same error levels under very general conditions [4]. In addition, it is easy to implement, reaches extremely high convergence rates and is completely independent of the underlying grid type (see Figure 1).

---

\*Correspondence to: [bmueLLer@gsc.tu-darmstadt.de](mailto:bmueLLer@gsc.tu-darmstadt.de)




Figure 1: Convergence study for the calculation of the arc-length of an ellipse

## References

- [1] J. Bremer, Z. Gimbutas, and V. Rokhlin. A nonlinear optimization procedure for generalized gaussian quadratures. *SISC*, 2010.
- [2] B. Engquist, A.-k. Tornberg, and R. Tsai. Discretization of dirac delta functions in level set methods. *J. Comput. Phys.*, 2004.
- [3] C. Min and F. Gibou. Geometric integration over irregular domains with application to level-set methods. *J. Comput. Phys.*, 2007.
- [4] B. Müller, F. Kummer, and M. Oberlack. Highly accurate surface and volume integration on implicit domains by means of moment-fitting. *Int. J. Numer. Meth. Engng.*, To be submitted.
- [5] B. Müller, F. Kummer, M. Oberlack, and Y. Wang. Simple multidimensional integration of smooth functions with discontinuous enrichments with application to level set methods. *Int. J. Numer. Meth. Engng.*, 2012.
- [6] S. E. Mousavi and N. Sukumar. Numerical integration of polynomials and discontinuous functions on irregular convex polygons and polyhedrons. *Comput. Mech.*, 2011.

## Acknowledgements

The work of B. Müller was supported by the ‘Excellence Initiative’ of the German Federal and State Governments and the Graduate School of Computational Engineering at Technische Universität Darmstadt.